

# Example Items

## Physics Pre-AP

**Physics Pre-AP Example Items** are a **representative set** of items for the ACP. Teachers may use this set of items along with the test blueprint as guides to prepare students for the ACP. On the last page, the correct answer, content SE and SE justification are listed for each item.

*The specific part of an SE that an Example Item measures is **NOT** necessarily the only part of the SE that is assessed on the ACP.* None of these Example Items will appear on the ACP.

Teachers may provide feedback regarding Example Items.

(1) Download the [Example Feedback Form](#) and email it. The form is located on the homepage of [Assessment.dallasisd.org](http://Assessment.dallasisd.org).

OR


(2) To submit directly, click “Example Feedback” **after** you login to the [Assessment website](#).

First Semester

2018–2019

Code #: 3211

# STAAR PHYSICS REFERENCE MATERIALS



## FORCE AND MOTION

$$\text{Average velocity} = \frac{\text{displacement}}{\text{change in time}}$$

$$v_{\text{avg}} = \frac{\Delta d}{\Delta t}$$

$$\text{Acceleration} = \frac{\text{final velocity} - \text{initial velocity}}{\text{change in time}}$$

$$a = \frac{v_f - v_i}{\Delta t}$$

$$\text{Acceleration} = \frac{(\text{final velocity})^2 - (\text{initial velocity})^2}{2(\text{displacement})}$$

$$a = \frac{v_f^2 - v_i^2}{2\Delta d}$$

$$\text{Displacement} = \left( \text{initial velocity} \right) \left( \text{change in time} \right) + \frac{1}{2} (\text{acceleration}) \left( \text{change in time} \right)^2$$

$$\Delta d = v_i \Delta t + \frac{1}{2} a \Delta t^2$$

$$\text{Centripetal acceleration} = \frac{(\text{tangential velocity})^2}{\text{radius}}$$

$$a_c = \frac{v_t^2}{r}$$

$$\text{Net force} = (\text{mass})(\text{acceleration})$$

$$F_{\text{net}} = ma$$

$$\text{Work} = (\text{force})(\text{distance})$$

$$W = Fd$$

$$\text{Torque} = (\text{force})(\text{lever arm})$$

$$\tau = Fr$$

$$\text{Power} = \frac{\text{work}}{\text{time}}$$

$$P = \frac{W}{t}$$

$$\text{Pythagorean theorem}$$

$$a^2 + b^2 = c^2$$

## GRAVITATIONAL, ELECTRICAL, AND MAGNETIC FORCES

$$\text{Force of gravitational attraction between 2 objects} = \left( \text{universal gravitation constant} \right) \left( \frac{\left( \text{mass of 1st object} \right) \left( \text{mass of 2nd object} \right)}{\left( \text{distance between centers of objects} \right)^2} \right)$$

$$F_g = G \left( \frac{m_1 m_2}{d^2} \right)$$

$$\text{Force between 2 charged particles} = \left( \text{Coulomb's constant} \right) \left( \frac{\left( \text{charge of 1st particle} \right) \left( \text{charge of 2nd particle} \right)}{\left( \text{distance between particles} \right)^2} \right)$$

$$F_{\text{electric}} = k_c \left( \frac{q_1 q_2}{d^2} \right)$$

$$\text{Electrical power} = (\text{voltage})(\text{current})$$

$$P = VI$$

$$\text{Current} = \frac{\text{voltage}}{\text{resistance}}$$

$$I = \frac{V}{R}$$


$$\text{Equivalent resistance for resistors in series}$$

$$R = R_1 + R_2 + R_3 + \dots$$

$$\text{Equivalent resistance for resistors in parallel}$$

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$$

# STAAR PHYSICS REFERENCE MATERIALS



## ENERGY AND MOMENTUM

$$\text{Kinetic energy} = \frac{1}{2}(\text{mass})(\text{velocity})^2 \qquad KE = \frac{1}{2}mv^2$$

$$\text{Gravitational potential energy} = (\text{mass})\left(\frac{\text{acceleration}}{\text{due to gravity}}\right)(\text{height}) \qquad PE_g = mgh$$

$$\text{Elastic potential energy} = \frac{1}{2}\left(\frac{\text{spring}}{\text{constant}}\right)\left(\frac{\text{distance stretched}}{\text{or compressed}}\right)^2 \qquad PE_{\text{elastic}} = \frac{1}{2}kx^2$$

$$\text{Energy} = (\text{power})(\text{time}) \qquad E = Pt$$

$$\text{Work} = \text{change in kinetic energy} \qquad W = \Delta KE$$

$$\text{Mechanical energy} = \text{kinetic energy} + \text{potential energy} \qquad ME = KE + PE$$

$$\text{Law of conservation of energy} \qquad KE_i + PE_i = KE_f + PE_f$$

$$\text{Momentum} = (\text{mass})(\text{velocity}) \qquad p = mv$$

$$\text{Impulse} = (\text{force})(\text{change in time}) = (\text{mass})(\text{change in velocity}) \qquad J = F\Delta t = m\Delta v$$

$$\text{Law of conservation of momentum} \qquad m_1v_{1i} + m_2v_{2i} = m_1v_{1f} + m_2v_{2f}$$

$$\text{Heat gained or lost} = (\text{mass})\left(\frac{\text{specific}}{\text{heat}}\right)\left(\frac{\text{change in}}{\text{temperature}}\right) \qquad Q = mc_p\Delta T$$


## WAVES AND LIGHT

$$\text{Velocity} = (\text{frequency})(\text{wavelength}) \qquad v = f\lambda$$

$$\frac{1}{\text{Focal length}} = \frac{1}{\text{distance to image}} + \frac{1}{\text{distance to object}} \qquad \frac{1}{f} = \frac{1}{d_i} + \frac{1}{d_o}$$

$$\text{Energy} = (\text{mass})(\text{speed of light})^2 \qquad E = mc^2$$

# STAAR PHYSICS REFERENCE MATERIALS



## CONSTANTS AND CONVERSIONS

$$c = \text{speed of light} = 3.00 \times 10^8 \frac{\text{m}}{\text{s}}$$

$$g = \text{acceleration due to gravity} = 9.8 \frac{\text{m}}{\text{s}^2}$$

$$G = \text{universal gravitation constant} = 6.67 \times 10^{-11} \frac{\text{N} \cdot \text{m}^2}{\text{kg}^2}$$

$$k_C = \text{Coulomb's constant} = 8.99 \times 10^9 \frac{\text{N} \cdot \text{m}^2}{\text{C}^2}$$

$$m_E = \text{mass of Earth} = 5.97 \times 10^{24} \text{ kg}$$

$$r_E = \text{radius of Earth} = 6.37 \times 10^6 \text{ m}$$

$$\text{newton (N)} = \frac{\text{kg} \cdot \text{m}}{\text{s}^2}$$

$$\text{joule (J)} = \text{N} \cdot \text{m}$$

$$\text{watt (W)} = \frac{\text{J}}{\text{s}} = \frac{\text{N} \cdot \text{m}}{\text{s}}$$

$$\text{hertz (Hz)} = \frac{\text{cycle}}{\text{s}}$$

# STAAR PHYSICS REFERENCE MATERIALS



1		2		3		4		5		6		7		8		9		10		11		12		13		14		15		16		17		18																																																																				
1A	1	2A	4	3B	4B	5B	6B	7B	8B	9B	10B	11B	12B	3A	4A	5A	6A	7A	8A	9A	10A	11A	12A	3A	4A	5A	6A	7A	8A	9A	10A	11A	12A																																																																					
	1 <b>H</b> 1.008 Hydrogen	2 <b>He</b> 4.003 Helium	3 <b>Li</b> 6.941 Lithium	4 <b>Be</b> 9.012 Beryllium	5 <b>B</b> 10.812 Boron	6 <b>C</b> 12.011 Carbon	7 <b>N</b> 14.007 Nitrogen	8 <b>O</b> 15.999 Oxygen	9 <b>F</b> 18.998 Fluorine	10 <b>Ne</b> 20.180 Neon	11 <b>Na</b> 22.990 Sodium	12 <b>Mg</b> 24.305 Magnesium	13 <b>Al</b> 26.982 Aluminum	14 <b>Si</b> 28.086 Silicon	15 <b>P</b> 30.974 Phosphorus	16 <b>S</b> 32.066 Sulfur	17 <b>Cl</b> 35.453 Chlorine	18 <b>Ar</b> 39.948 Argon	19 <b>K</b> 39.098 Potassium	20 <b>Ca</b> 40.078 Calcium	21 <b>Sc</b> 44.956 Scandium	22 <b>Ti</b> 47.867 Titanium	23 <b>V</b> 50.942 Vanadium	24 <b>Cr</b> 51.996 Chromium	25 <b>Mn</b> 54.938 Manganese	26 <b>Fe</b> 55.845 Iron	27 <b>Co</b> 58.933 Cobalt	28 <b>Ni</b> 58.693 Nickel	29 <b>Cu</b> 63.546 Copper	30 <b>Zn</b> 65.38 Zinc	31 <b>Ga</b> 69.723 Gallium	32 <b>Ge</b> 72.64 Germanium	33 <b>As</b> 74.922 Arsenic	34 <b>Se</b> 78.96 Selenium	35 <b>Br</b> 79.904 Bromine	36 <b>Kr</b> 83.798 Krypton	37 <b>Rb</b> 85.468 Rubidium	38 <b>Sr</b> 87.62 Strontium	39 <b>Y</b> 88.906 Yttrium	40 <b>Zr</b> 91.224 Zirconium	41 <b>Nb</b> 92.906 Niobium	42 <b>Mo</b> 95.96 Molybdenum	43 <b>Tc</b> (98) Technetium	44 <b>Ru</b> 101.07 Ruthenium	45 <b>Rh</b> 102.906 Rhodium	46 <b>Pd</b> 106.42 Palladium	47 <b>Ag</b> 107.868 Silver	48 <b>Cd</b> 112.412 Cadmium	49 <b>In</b> 114.818 Indium	50 <b>Sn</b> 118.711 Tin	51 <b>Sb</b> 121.760 Antimony	52 <b>Te</b> 127.60 Tellurium	53 <b>I</b> 126.904 Iodine	54 <b>Xe</b> 131.294 Xenon	55 <b>Cs</b> 132.905 Cesium	56 <b>Ba</b> 137.328 Barium	57 <b>La</b> 138.905 Lanthanum	58 <b>Ce</b> 140.116 Cerium	59 <b>Pr</b> 140.908 Praseodymium	60 <b>Nd</b> 144.242 Neodymium	61 <b>Pm</b> (145) Promethium	62 <b>Sm</b> 150.36 Samarium	63 <b>Eu</b> 151.964 Europium	64 <b>Gd</b> 157.25 Gadolinium	65 <b>Tb</b> 158.925 Terbium	66 <b>Dy</b> 162.500 Dysprosium	67 <b>Ho</b> 164.930 Holmium	68 <b>Er</b> 167.259 Erbium	69 <b>Tm</b> 168.934 Thulium	70 <b>Yb</b> 173.055 Ytterbium	71 <b>Lu</b> 174.967 Lutetium	72 <b>Hf</b> 178.49 Hafnium	73 <b>Ta</b> 180.948 Tantalum	74 <b>W</b> 183.84 Tungsten	75 <b>Re</b> 186.207 Rhenium	76 <b>Os</b> 190.23 Osmium	77 <b>Ir</b> 192.217 Iridium	78 <b>Pt</b> 195.085 Platinum	79 <b>Au</b> 196.967 Gold	80 <b>Hg</b> 200.59 Mercury	81 <b>Tl</b> 204.383 Thallium	82 <b>Pb</b> 207.2 Lead	83 <b>Bi</b> 208.980 Bismuth	84 <b>Po</b> (209) Polonium	85 <b>At</b> (210) Astatine	86 <b>Rn</b> (222) Radon	87 <b>Fr</b> (223) Francium	88 <b>Ra</b> (226) Radium	89 <b>Ac</b> (227) Actinium	90 <b>Th</b> 232.038 Thorium	91 <b>Pa</b> 231.036 Protactinium	92 <b>U</b> 238.029 Uranium	93 <b>Np</b> (237) Neptunium	94 <b>Pu</b> (244) Plutonium	95 <b>Am</b> (243) Americium	96 <b>Cm</b> (247) Curium	97 <b>Bk</b> (247) Berkelium	98 <b>Cf</b> (251) Californium	99 <b>Es</b> (252) Einsteinium	100 <b>Fm</b> (257) Fermium	101 <b>Md</b> (258) Mendelevium	102 <b>No</b> (259) Nobelium

Atomic number — 14 —  
Symbol — **Si** —  
Atomic mass — 28.086 —  
Name — Silicon —

Mass numbers in parentheses are those of the most stable or most common isotope.


Lanthanide Series		Actinide Series	
57 <b>La</b> 138.905 Lanthanum	58 <b>Ce</b> 140.116 Cerium	59 <b>Pr</b> 140.908 Praseodymium	60 <b>Nd</b> 144.242 Neodymium
89 <b>Ac</b> (227) Actinium	90 <b>Th</b> 232.038 Thorium	91 <b>Pa</b> 231.036 Protactinium	92 <b>U</b> 238.029 Uranium

## EXAMPLE ITEMS Physics Pre-AP, Sem 1

**1** John sees Hua running towards him at 11 m/s. While running, Hua throws a ball to John at 5 m/s. What is the speed of the ball as observed by John?

- A** 5 m/s
- B** 6 m/s
- C** 16 m/s
- D** 55 m/s

**2** A block with a mass of 5.0 kg is pushed on a frictionless surface through a distance of 5.0 m by applying a horizontal force of 80.0 N.




If the block starts from rest, what is its final velocity?

- A** 8.9 m/s
- B** 12.6 m/s
- C** 16.0 m/s
- D** 31.6 m/s

## EXAMPLE ITEMS Physics Pre-AP, Sem 1


3

An object moves in a straight line at a constant velocity of 1 m/s as shown in the velocity-time graph.




What position-time graph is represented by the motion indicated in the velocity-time graph?


A




C



B




D



## EXAMPLE ITEMS Physics Pre-AP, Sem 1

4

The diagram shows a 4.0 kg object accelerating at  $10 \text{ m/s}^2$  on a rough, horizontal surface.



What is the magnitude of the kinetic frictional force ( $F_{f, kinetic}$ ) acting on the object?

- A 50 N
- B 40 N
- C 20 N
- D 10 N

5

If the mass of a balloon is 300 kg and the lift force provided by the atmosphere is 3300 N, what is the net force on the balloon?


- A Upward, 360 N
- B Downward, 300 N
- C Toward the north, 300 N
- D Toward the east, 360 N

6

If the circles shown represent the relative sizes of four planets that all have the same mass, which planet exerts the greatest gravitational pull on objects on its surface?

A 

C 

B 

D 



## EXAMPLE ITEMS Physics Pre-AP, Sem 1

**7** A rock is thrown downward with a velocity of 25 m/s from the edge of a cliff which is 50 meters high. What is the approximate speed of the rock just before it hits the ground at the base of the cliff?

- A** 33 m/s
- B** 40 m/s
- C** 56 m/s
- D** 75 m/s

**8** The table shows the mass, velocity and height of four different objects.

Object	Mass (kg)	Velocity (m/s)	Height (m)
A	0.5	4.0	5
B	1.0	4.0	3
C	2.0	2.0	2
D	4.0	1.0	1

Which object has the most mechanical energy?

- A** Object A
- B** Object B
- C** Object C
- D** Object D


**9** Which statement accurately describes what happens as a ball falls freely (without air resistance) toward the ground?

- A** The total mechanical energy of the ball increases, but its kinetic energy decreases.
- B** The total mechanical energy of the ball decreases, but its potential energy increases.
- C** The total mechanical energy of the ball remains the same, but its kinetic energy decreases.
- D** The total mechanical energy of the ball remains the same, but its potential energy decreases.

## EXAMPLE ITEMS Physics Pre-AP, Sem 1

10

A rollercoaster cart is launched into a loop by a compressed spring that stores 10,000 Joules of energy.



Source: stickpng.com

If the starting position is considered zero height ( $h=0$ ) and there is no friction or air resistance, which pair of numbers describes possible values for the kinetic energy,  $KE$ , and the gravitational potential energy,  $PE$ , at point  $P$ ?

- A**      $KE = 0$  Joules  
           $PE = 11,000$  Joules
- B**      $KE = 9,000$  Joules  
           $PE = 3,000$  Joules
- C**      $KE = 3,000$  Joules  
           $PE = 7,000$  Joules
- D**      $KE = 7,000$  Joules  
           $PE = 1,000$  Joules

11

A toy car is rounding a circular track with a radius of 1.50 m. The car goes around the track once in 20.0 seconds. What is the centripetal acceleration of the car in  $\text{m/s}^2$ ?

(Circumference =  $2\pi r$ )

- A**      $0.12 \text{ m/s}^2$
- B**      $0.15 \text{ m/s}^2$
- C**      $0.31 \text{ m/s}^2$
- D**      $1.20 \text{ m/s}^2$

**EXAMPLE ITEMS Physics Pre-AP Key, Sem 1**

<b>Item#</b>	<b>Key</b>	<b>SE</b>	<b>Process Skills</b>	<b>SE Justification</b>
<b>1</b>	C	P.4B	2G, 2J	Describe motion relative to different frames of reference.
<b>2</b>	B	P.6A	2G, 2H, 2J	Calculate quantities using the work-energy theorem in various situations.
<b>3</b>	B	P.4A	2G, 2H, 2J	Interpret graphs describing different types of motion.
<b>4</b>	D	P.4D	2G, 2H, 2J	Calculate the effect of forces on objects, including the relationship between force and acceleration.
<b>5</b>	A	P.4D	2G, 2J	Calculate the effect of forces on objects.
<b>6</b>	A	P.5B	2H	Describe how the magnitude of the gravitational force between two objects depends on the distance between their centers.
<b>7</b>	B	P.4B	2G, 2J	Analyze motion in one dimension using equations with the concepts of displacement, instantaneous velocity, and acceleration.
<b>8</b>	C	P.6C	2H, 2J	Calculate the mechanical energy of a physical system.
<b>9</b>	D	P.6B	--	Investigate examples of kinetic and potential energy and their transformations.
<b>10</b>	C	P.6D	2G, 2J	Apply the law of conservation of energy.
<b>11</b>	B	P.4C	2G, 2H, 2J	Analyze accelerated motion in two dimensions using equations, including circular examples.